

PMRA APPROVED LABEL, JB
2020-09-22, Sub. No. 2020-0294

GROUP	4	HERBICIDE
--------------	----------	------------------

XtendiMax™ with VaporGrip™ Technology

HERBICIDE

SOLUTION

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: Dicamba, present as diglycolamine salt 350g a.e./L

REGISTRATION NO. 31896

PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 L to Bulk

READ THE LABEL AND ATTACHED BROCHURE BEFORE USING

KEEP OUT OF REACH OF CHILDREN

Bayer CropScience Inc.
Suite 200, 160 Quarry Park Blvd. SE
Calgary AB T2C 3G3
1-888-283-6847
www.cropscience.bayer.ca

RESTRICTED

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Thaw if frozen. Shake before use.

During mixing, loading, application, clean-up and repair, workers must wear long-sleeved shirt and long pants, chemical-resistant gloves and shoes plus socks. Chemical-resistant gloves are not required inside a closed cab or closed cockpit during application. For applications to non-crop areas, applicators must also wear coveralls

DO NOT enter treated fields until 12 hours after application to Roundup Ready 2 Xtend™ soybean and XtendFlex™ soybeans, barley, low bush blueberries, canary seed (*Phalaris canariensis*), corn (field and sweet), fallow, oats, pastures, red fescue, spring rye, seedling grasses, stubble fields, summer fallow and wheat (spring, durum).

Apply only when the potential for drift to areas of human habitation or activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

Surface Runoff

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to, heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted, fine textured or low in organic matter such as clay).

Potential contamination of aquatic areas as a result of runoff may be reduced by including an untreated vegetative strip between the treated area and the edge of the water body.

Avoid applying this product when heavy rain is forecast.

Leaching

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

FIRST AID

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

RESTRICTED

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For first aid instructions or the toxicological information essential for treatment, obtain and read the approved label from the registrant or phone the number indicated on this container.

TOXICOLOGICAL INFORMATION

Dicamba may cause severe irritation to the eyes and irritation to the skin and mucous membranes. Symptoms of overexposure to dicamba may include dizziness, muscle weakness, loss of appetite, weight loss, vomiting, decreased heart rate, shortness of breath, excitement, tenseness, depression, incontinence, cyanosis, muscle spasms, exhaustion and loss of voice. Treat symptomatically.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and it is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

STORAGE

- 1 Store **XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY HERBICIDE** in its original container only, away from other pesticides, fertilizer, food, or feed.
- 2 Keep the container closed to prevent spills and contamination.
- 3 Keep packages dry at all times.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

RESTRICTED

Product Specific Precautions

In case of an emergency involving this product, call Bayer CropScience day or night:
Accident/Spills/Medical Emergency 1-800-334-7577

For additional information on this or other Bayer CropScience agricultural products, call Bayer CropScience at: 1-888-283-6847

GROUP	4	HERBICIDE
--------------	----------	------------------

XtendiMax™ with VaporGrip™ Technology

HERBICIDE

SOLUTION

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: Dicamba, present as diglycolamine salt 350 g a.e./L

REGISTRATION NO. 31896 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 L to Bulk

READ THE LABEL AND ATTACHED BROCHURE BEFORE USING

KEEP OUT OF REACH OF CHILDREN

Bayer CropScience Inc.
Suite 200, 160 Quarry Park Blvd. SE
Calgary AB T2C 3G3
1-888-283-6847
www.cropscience.bayer.ca

RESTRICTED

ABOUT XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY HERBICIDE

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE controls broadleaf weeds in Roundup Ready 2 Xtend™ soybeans and XtendFlex™ soybeans, cereals, corn, reduced tillage (prior to seeding and reduced tillage fallow), pastures and rangeland grasses, crop-free land (summerfallow and stubble), red fescue, canary seed (*Phalaris canariensis*), seedling grasses grown for seed and forage, and low bush blueberries.

GENERAL PRECAUTIONS

1. **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** should not be applied on or near desirable trees or plants.
2. Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** when air temperature is between 10 and 25°C. Do not apply when there is a risk of severe fall in night temperature after use.
3. As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

4. Do not treat areas where movement of the chemical into the soil or surface washing may bring **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** into contact with roots of desirable plants.
5. Crop damage can occur if **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** is applied at any time other than the recommended crop stage.

NOTE: Crops growing under stress from adverse environmental conditions such as excess moisture, drought, disease, etc., may suffer a further setback and exhibit more pronounced injury symptoms if **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** is applied. However, the crop injury that may occur is usually offset by the weed control obtained.

6. Unless otherwise specified, do not use additives such as oil, wetting agents, emulsifiers, detergents, spreaders, sticking agents, or dispersing agents with **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** on crops.
7. If **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** is tank-mixed with another product, such as 2,4-D, consult that product's label for additional safety precautions, restrictions, application rates, timings and additional weeds controlled.
8. Ensure that spray equipment used to apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** is properly cleaned before re-using to apply any other chemicals. See section on suggested procedure for cleaning spray equipment.

RESTRICTED

SPRAY DRIFT PRECAUTIONS

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE may cause injury to desirable trees and plants, particularly non-Roundup Ready 2 Xtend™ Soybeans and XtendFlex™ Soybeans, flowers, fruit trees, grapes, ornamentals, peas, potatoes, tomatoes, tobacco, and other broadleaf plants especially in their developmental and growing stage. Follow these precautions when spraying in the vicinity of sensitive crops:

1. Treat when wind is 5 to 15 km/h. Do not apply during periods of dead calm or when weather conditions may cause drift from target areas to adjacent sensitive crops. Leave an adequate buffer zone between treatment areas and sensitive plants.
2. Use coarse sprays since they are less likely to drift than fine sprays. Select nozzles which minimize amounts of the fine spray particles. Do not exceed the nozzle manufacturer's recommended pressure, and use high flow rate (large orifice) nozzles to apply the highest practical spray volume.
3. Do not spray when the temperature is expected to exceed 30°C.
4. Avoid spraying under conditions of high humidity or fog.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

DIRECTIONS FOR USE

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) Coarse classification. Boom height must be 60 cm or less above the crop or ground.

Surface Runoff

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to, heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted, fine textured or low in organic matter such as clay).

Potential contamination of aquatic areas as a result of runoff may be reduced by including an untreated vegetative strip between the treated area and the edge of the water body.

Avoid applying this product when heavy rain is forecast.

Leaching

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

RESTRICTED

Buffer Zones

Use of the following spray methods or equipment DO NOT require a buffer zone: hand-held or backpack sprayer, spot treatment and inter-row hooded sprayer.

For application to rights-of-way, buffer zones for protection of sensitive terrestrial habitats are not required; however, the best available application strategies to minimize off-site drift, including meteorological conditions (e.g. wind direction, low wind speed) and spray equipment (e.g. coarse droplet sizes, minimizing height above canopy), should be used. Applicators must, however, observe the specified buffer zones for protection of sensitive aquatic habitats.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, rangelands, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

When a tank mixture is used, consult the labels of the tank mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

Buffer Zones Using ASAE Coarse Applications

Method of Application	Crop	Buffer Zones (meters) Required for Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field Sprayer	Roundup Ready 2 Xtend™ soybean and XtendFlex™ soybeans, Field Corn, Summer fallow and stubble fields, Perennial rosette in summer fallow field, Reduced tillage fallow field, New/established red fescue for seed, Established grass pasture	1	1	0	0	4
	Cereals (spring wheat, spring barley, spring rye, winter wheat, oats), canary grass (seed production), Seed and forage crop production (brome grass, smooth fescue, meadow fescue, tall foxtail, meadow orchard grass, red fescue, creeping timothy and wheatgrass (crested, intermediate, pubescent, slender, streambank and tall).	0	0	0	0	1

	Reduced tillage prior to seeding (in wheat, barley, rye, oats, and corn (except sweet corn))	0	0	0	0	4
	Pasture and rangeland, non-cropland**	1	1	0	0	10
	Lowbush blueberries	1	1	1	0	15
Field Sprayer	Roundup Ready 2 Xtend and XtendFlex™ Soybeans Seed Production	1	1	0	0	10

** Buffer zones for the protection of terrestrial habitats are not required for use on rights-of-way, including railroad ballast, rail and hydro rights-of-way, utility easements, roads, and training grounds and firing ranges on military bases.

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pesticides portion of the Canada.ca website

DIRECTIONS FOR USE

ROUNDUP READY 2 XTEND™ SOYBEANS and XTENDFLEX™ SOYBEANS

XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide and Roundup WeatherMAX® with Transorb® 2 Technology Liquid Herbicide Use In Roundup Ready 2 Xtend Soybeans and XtendFlex™ Soybeans

Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** only to Roundup Ready 2 Xtend™ Soybeans and XtendFlex™ Soybeans. Roundup Ready 2 Xtend™ Soybean and XtendFlex™ Soybeans contains a patented gene that provides tolerance to dicamba, the active ingredient in this product. This product will cause severe crop injury or destruction and yield loss if applied to soybeans that are not dicamba tolerant, including soybeans with a trait engineered to confer tolerance to auxin herbicides other than dicamba.

Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** alone or in tank mixture with Roundup WeatherMAX with Transorb 2 Technology Liquid Herbicide in 100-220 L/ha

DO NOT apply using aerial application equipment.

RESTRICTED

Weeds Controlled *	Rates	Timing
<p>Annual Broadleaf Weeds: biennial wormwood (1) buckwheat (wild, tartary) bur cucumber (2) Canada fleabane chickweed cleavers cocklebur common lamb's-quarters corn spurry cow cockle Eastern black nightshade flixweed kochia lady's-thumb mustard (hare's-ear, Indian, tumble, wild, wormseed) narrow leaved hawk's-beard night flowering catchfly non-glyphosate tolerant canola (rapeseed) hempnettle pigweed (redroot, Russian, smooth) ragweed (common, false, giant) Russian thistle shepherd's-purse smartweed (green, Pennsylvania) stinkweed stork's bill velvetleaf volunteer adzuki beans (3) wild tomato</p> <p>Annual Grass Weeds: barnyard grass crabgrass (smooth, large) fall panicum foxtail (green, yellow, giant) volunteer barley volunteer wheat wild oats wild proso millet</p> <p>Perennial Weeds: Canada Thistle(4) common milkweed(5,7) dandelion (6) field bindweed(7) perennial sow thistle(4) quackgrass(4)</p>	<p>XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 823 mL to 1.71 L/ha + Roundup WeatherMAX at 1.67 L/ha</p>	<p>Preplant or Preemergence to the crop and/or Postemergence to the crop once or twice up to the early flower stage of the crop (R1)</p> <p>Notes: The 1.71 L/ha rate of XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE is to be used only once in a season and should be used preplant, preemergence or in-crop early postemergence</p> <p>3.36 L/ha of XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE is the maximum total to be applied to Roundup Ready 2 Xtend Soybeans and XtendFlex™ Soybeans in a single growing season (year).</p> <p>A third application of of XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE should only be made for the control of glyphosate resistant weed populations.</p> <p>Also see Residual Weed Control Section below for more information.</p>

wire-stemmed muhly(4) yellow nutsedge (5,7),		
All Weeds listed above plus tall water hemp (8), and horsenettle (8),	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 823 mL to 1.71 L/ha + Roundup WeatherMAX at 3.33 L/ha	See notes above for application details. - 1 application per season at 3.33 L/ha.
All Weeds listed above plus volunteer alfalfa (9) and bromegrass (9)	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 823 mL to 1.71 L/ha + Roundup WeatherMAX at 4.67 L/ha (10)	See notes above for application details. - 1 application per season at 4.67 L/ha.

* Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of annual weeds greater than 25 cm in height will be inconsistent, although some weeds may be controlled.

Application Footnotes:

(1) One application including Roundup WeatherMAX at 1.67 L/ha applied at the 2-8 leaf stage of actively growing biennial wormwood.

(2) Two applications including Roundup WeatherMAX at 1.67 L/ha applied when the bur cucumber is at the 1 to 18 leaf stage. Applications should be at least 2 weeks apart for best results.

(3) Applications including Roundup WeatherMAX at 1.67 L/ha applied at the unifoliate to 4th trifoliate leaf stage of the Adzuki beans. A second 1.67 L/ha application may be used for late flushes emerging after the initial treatment when the Adzuki beans are in the unifoliate to 4th trifoliate leaf stage and actively growing.

(4) Applications including Roundup WeatherMAX at 1.67 L/ha applied when quackgrass has 3-4 leaves, Canada thistle and perennial sow thistle are rosette to 50 cm in height, and wire-stemmed muhly is 10-20 cm in height. Weeds should be actively growing at application.

(5) Applications including Roundup WeatherMAX at 1.67 L/ha will provide suppression.

(6) Applications including Roundup WeatherMAX applied preplant surface or pre-emergence at 1.67 to 3.33 L/ha. Use Roundup WeatherMAX rates of 2.47 to 3.33 L/ha on heavy infestations of dandelions and on dandelions greater than 15 cm in size. Apply up to and including bloom for best results.

(7) For control of common milkweed, yellow nutsedge, and field bindweed a second application including Roundup WeatherMAX at 1.67 L/ha may be needed and should be applied at least 2 weeks after the first application or 3.33 L/ha should be applied once. Milkweed should be 15-60 cm in height, yellow nutsedge should be 5-15 cm in height.

(8) Applications including Roundup WeatherMAX at 3.33 L/ha applied at the 2-12 leaf stage of horse-nettle or up to the 18 leaf stage of tall waterhemp or 2 applications of 1.67 L/ha applied at least 2 weeks apart. For control of tall waterhemp use the higher rate if weeds are beyond the 6 leaf stage.

(9) Alfalfa should have 9 or more leaves and be at least 10-15 cm tall. Bromegrass should have at least 3-5 leaves and be at least 10-15 cm tall.

(10) With the 4.67 L/ha rate some short term yellowing may occur in the sprayer overlap areas, but this effect is temporary and will not influence growth or yield.

Residual Weed Control and Suppression with XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Applications

In addition to providing postemergence burndown activity on weeds **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** applications will also provide short term residual activity on the weeds listed below.

Residual Weed Control and Suppression provided with XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Applications (the 1.71 L/ha rate provides short term control and the 823 mL/ha rate provides suppression)
common lamb's-quarters common ragweed redroot pigweed velvetleaf* wild buckwheat *suppression only for both rates

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Use In Roundup Ready 2 Xtend™ Soybeans and XtendFlex™ Soybeans

Weeds Controlled	Rates	Timing
<p>Annual Broadleaved Weeds: buckwheat (Tartary, wild) Canada fleabane (1) cleavers common lamb's-quarters corn spurry cow cockle green smartweed lady's-thumb, mustard (hare's-ear, Indian, tumble, wild, wormseed) pigweed (redroot, Russian, smooth) ragweed (common, false, giant) velvetleaf</p> <p>Perennial Weeds: Canada Thistle (2) field bindweed (2) perennial sow thistle (2)</p>	<p>XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 823 mL to 1.71 L/ha</p>	<p>Preplant or Preemergence to the crop</p> <p>and/or</p> <p>Postemergence to the crop once or twice up to the early flower stage of the crop (R1)</p> <p>Notes: The 1.71 L/ha rate of XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE is to be used only once in a season and should be used preplant, preemergence or in-crop early postemergence</p> <p>3.36 L/ha of XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE is the maximum total to be applied to Roundup Ready 2 Xtend Soybeans and XtendFlex™ Soybeans in a single growing season (year).</p> <p>A third application of XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE, should only be made for the control of glyphosate resistant weed populations.</p>

Application Footnotes :

- (1) Post-emergence application only
- (2) Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** annually for three years at the flowering stage of bindweed and the budding stage of thistles.

Pre-Harvest Interval(s):

7-10 days for soybean forage and 13-15 days for soybean hay.

Rotational Crop Guidelines:

A plant back interval of 120 days is required for those crops not on the **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** label.

Do not count days when the ground is frozen. Moisture is essential for the degradation of this herbicide in soil. If dry weather persists after application, crop injury may occur the following spring.

Equipment – DO NOT APPLY THIS PRODUCT TO ROUNDUP READY 2 XTEND SOYBEANS AND XTENDFLEX SOYBEANS USING AERIAL SPRAY EQUIPMENT.

Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** to weeds < 10 cm.

Water Volume – apply this product in a minimum of 100 Liters of spray solution per hectare.

Spray Drift Management

Do not allow herbicide solution to mist, drip, drift or splash onto desirable vegetation because severe injury or destruction to desirable broadleaf plants could result. The following drift management requirements must be followed to ensure application accuracy from ground application onto agricultural field crops.

Controlling Droplet Size

The most effective way to reduce spray drift potential is to apply large droplets (coarser spray qualities) that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if the application is made improperly, or under unfavorable environmental conditions (see the “**Wind Speed and Direction**”, “**Temperature and Humidity**” and “**Temperature Inversions**” sections of this label).

- **Nozzle type.** Use only spray nozzles that produce Extremely Coarse to Ultra Coarse spray qualities and minimal amounts of fine spray droplets as defined by the American Society of Agricultural Engineers (ASAE S-572.1). Do not use conventional flat fan nozzles that produce Medium or Fine spray qualities.
Check nozzle manufacturer’s recommendations to determine the proper operating pressure, nozzle spacing and ground speed that will deliver Extremely Coarse to Ultra Coarse spray qualities at spray volumes of at least 100 L/ha.
- **Spray Pressure.** Adjust pressure for selected nozzles according to the nozzle manufacturer to maintain Extremely Coarse to Ultra Coarse spray qualities. Do not exceed the nozzle manufacturer’s recommended pressures. Use sufficient spray pressure with air induction nozzles to ensure a good spray pattern, while maintaining Extremely Coarse to Ultra Coarse sprays; use at least 200 kPa to ensure proper pattern overlap and check this visually. Confirm that sprayer rate controller hardware (if so equipped) does not increase pressure above the range that produces the correct spray quality. Calibrate the flow rate for the selected nozzles on the equipment used to apply this product.
- **Spray Volume.** Apply this product in a minimum of 100 Liters of spray solution per hectare. Use a higher spray volume when treating dense vegetation. Higher spray volumes also allow the use of larger nozzle orifices (sizes) which produce coarser spray droplets.

- **Equipment Ground Speed.** Select a ground speed under 25 km/h that will deliver the desired spray volume while maintaining the desired spray pressure. Slower speeds generally result in more uniform spray coverage and deposition on the target area.

- **Spray boom Height.** Spray at the appropriate boom height based on nozzle selection and nozzle spacing (should not be more than 50 cm above target pest or crop canopy). Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions. Automated boom height controllers are recommended with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions.

Do not apply during a temperature inversion because off-target movement potential is high.

- During a temperature inversion, the atmosphere is very stable and vertical air mixing is restricted, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions.
- Temperature inversions are characterized by increasing temperatures with altitude and are common on evenings, nights, and early mornings with limited cloud cover and light to no wind. After sunset, air at the earth's surface cools and air is trapped by warmer air above it. Inversions begin to form as the sun sets and often continue into the morning.
- Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.
- The inversion will dissipate with increased winds (above 5 km/h) or after sunrise when the surface air begins to warm.

Wind Speed and Direction

- Drift potential is lowest between wind speeds of 5 to 15 km/h.
- If the wind speed is 5 km/h or less and fog is present, indicating a temperature inversion, do not apply this product.
- If fog is not present, conduct a smoke test. Smoke that moves upward confirms there is no inversion present whereas smoke that layers and moves laterally in a concentrated cloud, indicates a temperature inversion exists. Do not apply this product during a temperature inversion. Wait until the wind speed is greater than 5 km/h to ensure that any inversion has lifted.
- Do not spray this product when the wind is blowing in the direction of a sensitive area at a wind speed greater than 15 km/h.
- For **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** wind speed and direction restrictions see below table:

Wind speed	Application conditions and restrictions
<5 km/h	Do not apply XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE if temperature inversion exists.
5-15 km/h	Optimum XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE application conditions.
>15 km/h	DO NOT apply XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE .

NOTE: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

Additives and tank mixes

- Nozzle selection is one of the most important parameters for drift reduction. A drift reduction additive may be used with this product to further reduce fine droplets.
- Not all drift reduction additives are compatible with every nozzle type and pesticide / adjuvant combination. Check with the additive manufacturer to insure that the drift additive will work properly with the spray nozzle, spray pressure and your specific spray solution. Read and carefully observe all precautions, limitations and all other information on the product label.
- A quality nonionic surfactant (NIS) of at least 70% active may be added to the spray solution at 0.25 % v/v. Read and carefully observe all caution statements and other information on the surfactant label.
- Do not add acidifying buffering agents, acidic pH adjusting agents or adjuvants other than agriculturally approved NIS to the spray solution. Do not add ammonium sulfate (AMS), AMS-containing adjuvants, water conditioners, or sprayable fluid fertilizers.
- Do not use crop oil concentrates (COC) and methylated seed oils (MSO) as adjuvants when this product is applied with glyphosate-based agricultural herbicides. When **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** is used with another herbicide that requires the use of a COC or MSO adjuvant follow the label instructions of that product.
- In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Bayer CropScience at 1-888-283-6847 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.
- Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** or tank mixtures with **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** at a minimum spray volume of 100 L/ha.

Sensitive Areas

This product should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for species at risk, or sensitive crop plants) is minimal (e.g. when the wind is blowing away from sensitive areas). Applicators should survey the surrounding area before making an application of this product.

Failure to follow the requirements in this label, could result in severe injury or destruction to desirable sensitive crops and trees, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes, and other broadleaf plants when contacting their roots, stems or foliage.

Application Awareness

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of equipment and weather related factors must be monitored to maximize performance and on-target spray deposition. The applicator is responsible for considering all of these factors when making a spray decision.

Proper spray system equipment cleanout

Minute quantities of dicamba may cause injury to **non-Roundup Ready 2 Xtend soybeans and XtendFlex soybeans** and other sensitive crops (see the “Sensitive Areas” section of this label for a listing of sensitive crops).

Clean equipment immediately after using this product, using a triple rinse procedure as follows:

1. After spraying, drain the sprayer (including boom and lines) immediately. Do not allow the spray solution to remain in the spray boom lines overnight prior to flushing.
2. Flush tank, hoses, boom and nozzles with clean water.
3. Inspect and clean all strainers, screens and filters.
4. Prepare a cleaning solution with a commercial detergent or sprayer cleaner or ammonia according to the manufacturer’s directions.
5. Take care to wash all parts of the tank, including the inside top surface. Start agitation in the sprayer and thoroughly re-circulate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.
6. Flush hoses, spray lines and nozzles for at least 1 minute with the cleaning solution.
7. Repeat above steps for two additional times to accomplish an effective triple rinse.
8. Remove nozzles, screens and strainers and clean separately in the cleaning solution after completing the above procedures.
9. Appropriately dispose of rinsate from steps 1-7 in compliance with all applicable laws and regulations.
10. Drain sump, filter and lines.
11. Rinse the complete spraying system with clean water.

All rinse water must be disposed of in compliance with municipal, provincial, and federal guidelines.

Seed Production in ROUNDUP READY 2 XTEND SOYBEANS and XTENDFLEX™ SOYBEANS

Weeds Controlled	Rates and Timing	
<p>Non tolerant (off type) Roundup Ready 2 Xtend soybean lines</p>	<p>Treatment Option 1:</p> <p>XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY HERBICIDE at 3.36 L/ha - Late postemergent (soy V6-R1)</p> <p>OR</p> <p>Treatment Option 2:</p> <p>XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY HERBICIDE (1.71 L/ha) - Early post emergent (soy V1-V3) followed by XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY HERBICIDE (3.36 L/ha) - Late post emergence (soy V6-R1)</p>	<p>DO NOT exceed 5.14 L/ha of XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY HERBICIDE per year.</p>

Pre-Harvest Interval(s): 86 days for soybean seed, 7 days for soybean forage and 14 days for soybean hay.

CEREALS (not underseeded to legumes)

Treatment Notes

1. For best performance, spray when weeds are in the 2 to 3 leaf stage and rosettes are less than 5 cm across.
2. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.
3. Crop damage can occur if application is made at any time other than the recommended crop stage.
4. Do not apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** or **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mixes if crop is under-seeded to legumes.

Application Directions

Ground Application

Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** or **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mixes in at least 110 litres of water/ha.

Application of this specific product must meet and/or conform to the following:

- 1.
2. Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** alone at 230 mL/ha or tank mix **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** at 230 mL/ha with the recommended rate of the phenoxy herbicides specified on this label.
3. Treat when wind is 5 to 15 km/h. Do not apply during periods of dead calm or when weather conditions may cause drift from target areas to adjacent sensitive crops.
4. Do not use nozzle pressure above 200 kPa.
5. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, or shelterbelt.
6. Unless otherwise specified, do not use any additives with **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE**.

Weeds Controlled

Weeds Controlled	Rates	Tank Mix
Canada thistle (top growth only) cleavers (higher rate only) corn spurry cow cockle green smartweed lady's thumb perennial sow-thistle (top growth only), Tartary Buckwheat wild buckwheat,	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone at 315-398 mL/ha	None
Weeds listed for XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone plus: burdock (young seedlings) cocklebur flixweed hemp-nettle** kochia redroot pigweed Russian pigweed Russian thistle shepherd's-purse volunteer canola* volunteer sunflower*** wild radish	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 315 mL/ha +	2, 4-D amine OR MCPA amine OR MCPA K

Weeds listed for XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone plus: chickweed hemp-nettle** stinkweed volunteer sunflower***	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 315 mL/ha +	Sencor OR Lexone
Weeds listed for XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone plus: volunteer canola*	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 315 mL/ha +	Ally

* Best results will be obtained if application is made prior to bolting of canola, when this weed is at the 2 to 4 leaf stage.

** Use **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** + MCPA K for hemp-nettle control. Apply at the 2 to 3 leaf stage of weed for best control. Hemp-nettle may not be controlled if application is made at a more advanced stage of crops and weeds.

*** Depending on the growing conditions, control may be slightly delayed.

Application Directions

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE may be applied to:

- Spring Wheat
- Spring Barley
- Winter Wheat
- Oats
- Spring Rye

The following sections describe application directions for these crops.

Spring Wheat

Herbicide	Rate	Crop Stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone	315-398 mL/ha	2-5 leaf
+ 2,4-D amine	850 mL/ha (500 g/L formulation)	2-5 leaf
or MCPA amine	850 mL/ha (500 g/L formulation)	2-5 leaf
or MCPA K	1.1 L/ha (400 g/L formulation)	2-5 leaf
or Sencor 500*	275-425 mL/ha**	2-3 leaf
or Lexone DF*	275 g/ha	2-3 leaf
or Ally***	5 g/ha	2-5 leaf

* Sencor/Lexone tank-mixes apply to Western Canada only. Application may be delayed until the 4-leaf stage of the crop, however, crop tolerance may be reduced. Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** at 315 mL/ha with Sencor/Lexone.

** Use the higher rate of Sencor 500 for control of volunteer sunflowers.

*** Ally tank-mixes apply to Western Canada only. Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** at 315 mL/ha with Ally. Ensure that Ally is completely in suspension in the spray tank before adding **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE**. Do not add a surfactant.

Spring Rye

Herbicide	Rate	Crop Stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone	315-398 mL/ha	2-3 leaf
+ 2,4-D amine	850 mL/ha (500 g/L formulation)	2-3 leaf

Spring Barley

Herbicide	Rate	Crop Stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone	315-398 mL/ha	2-5 leaf
+2,4-D amine	850 mL/ha (500 g/L formulation)	2-5 leaf
or MCPA amine	850 mL/ha (500 g/L formulation)	2-5 leaf
or MCPA K	1.1 L/ha (400 g/L formulation)	2-5 leaf
or Sencor 500*	275-425 mL/ha**	2-3 leaf
or Lexone DF*	275 g/ha	2-3 leaf
or Ally***	5 g/ha	2-5 leaf

* Sencor/Lexone tank-mixes apply to Western Canada only. NOTE: Do not use on Klondike barley.

** Use the higher rate of Sencor 500 for control of volunteer sunflowers.

*** Ally tank-mixes apply to Western Canada only. Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** at 315 mL/ha with Ally. Ensure that Ally is completely in suspension in the spray tank before adding **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE**. Do not add a surfactant.

Winter Wheat

Herbicide	Rate	Crop Stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone	315-398 mL/ha	15-25 cm tall or before shot-blade stage
+ 2,4-D amine	850 mL/ha (500 g/L formulation)	15-25 cm tall or before shot-blade stage
or MCPA amine	850 mL/ha (500 g/L formulation)	
or MCPA K	1.1 L/ha (400 g/L formulation)	

Oats

Herbicide	Rate	Crop Stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone	315-398 mL/ha	2-5 leaf
+ MCPA amine	850 mL/ha (500 g/L formulation)	2-5 leaf
or MCPA K	1.1 L/ha (400 g/L formulation)	2-5 leaf

Grazing Restrictions:

Following treatment with **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** or **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** plus 2,4-D, follow these grazing restrictions:

- DO NOT permit lactating dairy animals to graze fields within 7 days after application.
- DO NOT harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

Following treatment with **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** plus any other herbicide tank mix: Do not graze or harvest for livestock feed prior to crop maturity; sufficient data are not available to support such use.

FIELD CORN

DO NOT apply using aerial application equipment.

Treatment Notes

1. Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** or **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mixes in 220 to 350 litres of water/ha at a pressure of 150 to 275 kPa. Use coarse sprays.
2. Keep spray mixture in suspension at all times. If mixture is allowed to settle, thoroughly agitate the mixture before spraying.
3. Do not apply to sweet corn.
4. Unless otherwise specified, do not use additives such as oil, wetting agents, emulsifiers, detergents, spreaders, sticking agents, or dispersing agents on corn with **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE**.
5. Corn height refers to the crop as it stands, not leaf-extended.
6. When using drop pipes (drop nozzles), direct the spray beneath the lower leaves of the corn and onto the weeds and soil. Do not apply to corn over 50 cm in height.
7. Apply no later than 2 weeks prior to tassel emergence when using **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** alone up to 50 cm.
8. For the best control of annuals, spray when they are actively growing and in the seedling stage. Poor results may occur if weeds are well advanced at the time of application.
9. When applying **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** adjacent to sensitive crops, apply as a pre-emergent or early post-emergent treatment to avoid potential drift onto these sensitive crops.
10. When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions and grazing limitations for each product used in the tank-mix. Follow the more stringent label precautionary and PPE measures for mixing/loading/applying, and label statements pertaining to environmental protection, such as buffer zones, stated on all tank-mix product labels.

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE / LIQUID NITROGEN

Pre-emergent applications of **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** are generally compatible with most liquid nitrogen fertilizers. To determine compatibility, mix all components of the finished spray in proportionate quantities in a small jar before mixing in the spray tank. If the herbicides do not ball-up or form flakes, sludge, jelly, oily films or layers, or other precipitates within 5 minutes after mixing, the tested spray-mix is compatible.

Weeds Controlled

Weeds Controlled	Rate	Crop Stage
buckwheat (tartary, wild), Canada fleabane*** Canada thistle** cleavers common ragweed** corn spurry cow cockle false ragweed field bindweed ** giant ragweed green smartweed lady's-thumb lamb's-quarters* mustard, (hare's-ear, Indian, tumble, wild, wormseed) perennial sow-thistle** redroot pigweed** Russian pigweed velvetleaf	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone at 823 mL – 1.71 L/ha	none

* Including atrazine resistant species

** Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** annually for three years at the flowering stage of bindweed and the budding stage of thistles.

*** Post emergent applications only

Pre-Emergence Treatment

Eastern Canada Only

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE can be used alone at 1.71 L/ha or in tank-mixes with the following herbicides for additional broadleaf and grassy weed control.

Herbicide	Rate L/ha
Dual Magnum	1.25 – 1.75L
Dual II Magnum	1.25 – 1.75 L
Frontier Max Herbicide	0.756 – 0.963 L
Primextra II Magnum	3.0 - 4.0 L
Aatrex Liquid 480*	2.10 – 3.10L
Aatrex Liquid 480* + Dual II Magnum	2.10 – 3.10 L + 1.25 – 1.75 L

* Other atrazine formulations will require a rate calculation adjustment according to percent active ingredient

Pre-Emergence Treatment Notes

- Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mixes as broadcast ground treatments after planting but before weeds and corn emerge.
- Apply to medium to fine textured soils containing more than 2.5% organic matter.
- Do not use on sandy or sandy loam soils.
- Avoid direct chemical contact with the corn seed. If you plan to apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** prior to corn emergence, be sure to place the corn seeds 4 cm or more below the soil surface. If seeds are planted less than 4 cm below the soil surface, delay application of **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** until the spike stage.
- Do not incorporate. If applications are made during planting, apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** far enough behind the planting equipment to avoid incorporation by the planter wheel or other covering device. If soil crusting makes it necessary to use a rotary hoe after a pre-emergence treatment, delay hoeing the soil more than 1.3 cm deep.
- Always consult the tank mix partner label for further limitations and restrictions (especially re: soil type).

Post-Emergence Treatment

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE or **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mixes can be applied as “overlay” to corn previously treated with any other broadleaf or grass herbicide. The 1.71 L/ha rate of **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** as “overlay” is particularly effective in controlling velvetleaf and providing extended residual control of other late germinating, deep rooted annuals. **Note:** Unless otherwise specified, do not use additives such as oils, wetting agents, or sticking agents.

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone**Spike to 5-leaf corn****Eastern and Western Canada**

Herbicide	Rate	Crop Stage	Weed Stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone	1.71 L/ha	Spike to 5-leaf	Pre-emergence to 2-leaf ¹

¹ For best performance, spray when the broadleaf weeds are emerged and up to the 2-leaf stage of their development.

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Tank-mixes**Western Canada (Prairie Provinces only)***

Herbicide	Rate	Crop Stage	Weed Stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE + Accent 75DF + non-ionic surfactant such as Agral®, Agsurf® or Citowett® Plus	0.823 L (288 g ai/ha) + 33 g (25 g ai/ha) + 0.2% v/v	Spike to 6-leaf	Post-emergence to 6-leaf

* Single post-emergent spray; ground application only; do not apply this tank mix within 30 days of harvest.

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE tank-mixes
Eastern Canada only

Herbicide	Rate/ha	Crop Stage	Weed Stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE + Frontier Max Herbicide	1.71 L + 0.756 – 0.963 L	Spike to 3-leaf	Pre-emergence to 2-leaf**
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE + Aatrex Liquid 480*	1.71 L + 2.10 – 3.10 L	Spike to 5-leaf	Pre-emergence to 2-leaf
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE+ Aatrex Liquid 480* + Dual II Magnum	0.823 - 1.71 L + 2.10 – 3.10 L + 1.25 – 1.75 L	Spike to 2-leaf	Emergence to 2-leaf
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE + Primextra II Magnum	0.823 -1.71 L + 3.0 - 4.0 L	Spike to 2-leaf	Emergence to 2-leaf
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE + Ultim 75% DF + non-ionic surfactant	0.823 L + 1 bag + 0.2% v/v	Spike to 6-leaf	Emergence to 6-leaf
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE + Elim EP Herbicide 25% DF + non-ionic surfactant	0.823 L + 60 g + 0.2% v/v	Spike to 3-leaf	Emergence to 4-leaf
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE + Dual II Magnum	0.823 -1.71 L + 1.25 - 1.75 L	Spike to 2-leaf	Emergence to 2-leaf

* Other atrazine formulations will require a rate calculation adjustment according to percent active ingredient

** For annuals, apply before 2-leaf stage.

**XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE tank-mixes
Eastern Canada and the Province of Manitoba***

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE can be tank mixed with Option 35 DF herbicide and applied as a post-emergence application to field corn grown in Eastern Canada and the province of Manitoba. Tank mixing **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** with Option 35 DF will provide enhanced control of annual broadleaf weeds.

Option 35 DF herbicide is to be used in conjunction with Hasten spray additive at 1.75 L/ha plus liquid nitrogen fertilizer (28% UAN) at a rate of 2.5 L/ha. Use of a spray-grade liquid nitrogen fertilizer is recommended.

Herbicide	Rate/ha	Corn Stage	Weed Stage	Weeds Controlled
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE+ Option 35 DF + Hasten spray adjuvant + liquid nitrogen fertilizer (28% UAN)	0.4 L + 100 g + 1.75 L + 2.5 L	1 to 8-leaf	Consult the Option 35 DF label for the recommended leaf stage of weeds at application. For best results, apply to emerged, young, actively growing weeds.	<p>Perennials quackgrass</p> <p>Annual Grasses barnyard grass bristly foxtail fall panicum green foxtail large crab grass proso millet witchgrass yellow foxtail</p> <p>Annual Broadleaf Weeds common chickweed common ragweed (suppression only) Eastern black nightshade lamb's-quarters redroot pigweed velvetleaf wild mustard wormseed mustard</p>

*Ground application only. Do not apply by air. Make only one application per season. Apply in a minimum of 220 L/ha of water. and at a pressure of 175 – 275 kPa.

**Spike to 50 cm standing corn
Eastern and Western Canada**

Herbicide	Rate/ha	Corn Stage	Weed Stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone	823 mL	Emergence to 50 cm (drop nozzles from 20 - 50 cm corn)	Pre-emergence to 2-leaf
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE + 2,4-D amine	398 mL + 850 mL	Emergence to 50 cm (drop nozzles from 20 - 50 cm corn)	Pre-emergence to 2-leaf

**Sequential XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Applications
Eastern and Western Canada**

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE may be applied sequentially to a **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** application to control late-emerging weeds such as field bindweed, Canada thistle and velvetleaf. Follow application directions as outlined for the **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** alone post-emergence treatments up to 50 cm tall corn.

Grazing Restrictions:

DO NOT permit lactating dairy animals to graze fields within 7 days after application

DO NOT harvest forage or cut hay within 30 days after application

Withdraw meat animals from treated fields at least 3 days before slaughter

WEED CONTROL IN REDUCED TILLAGE (prior to seeding)

DO NOT apply using aerial application equipment.

Treatment Notes

1. **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE + Roundup brand** agricultural products (glyphosate only) applications may be applied to emerged annual grass and annual broadleaf weeds in reduced tillage systems prior to seeding of spring wheat, spring barley spring rye, winter wheat, oats, and field corn only.
2. Do not apply prior to seeding sweet corn.
3. Planting should follow soon after application since this tank-mix does not provide residual weed control.
4. Delayed planting following chemical application will allow weeds to emerge between application and crop emergence.
5. For field corn, apply to medium to fine textured soils containing more than 2.5% organic matter. Do not use on sandy or sandy loam soil.
6. Certain broadleaf crops such as sweet corn, lentils, peas, canola and flax can be injured by a pre-seeding application of this tank-mix and should not be planted after the use of this tank-mix.

7. Under certain stress conditions, such as drought, cool temperatures or where extremely hard water (> 700 ppm Ca + Mg) will be used, use 50 L/ha of water with this tank-mix to help improve results. However, the combination of low water volume and coarse sprays may reduce spray coverage.

Application Directions

Weeds Controlled	Rate	Tank Mix
Annual Grasses (Apply any time between emergence and heading): downy brome green foxtail Persian darnel volunteer cereals wild oats	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 432 mL/ha +	330-686 g ae/ha glyphosate+ 0.5 L of a non-ionic surfactant in 100 L of water
Annual Broadleaves (Apply up to 15 cm height): cleavers (1-4 whorls) (suppression only) cow cockle flixweed** green smartweed kochia lady's-thumb lamb's-quarters redroot pigweed Russian thistle stinkweed** volunteer canola*** wild buckwheat* wild mustard	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 432 mL/ha +	330-686 g ae/ha glyphosate+ 0.5 L of a non-ionic surfactant in 100 L of water
Perennials (Apply before initiation of seed head or browning of lower leaves): foxtail barley (suppression only)	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 432 mL/ha +	glyphosate 330-686 g ae/ha glyphosate + 0.5 L of a non-ionic surfactant in 100 L of water

* Apply at the 1 to 4-leaf stage.

** For optimal control of winter annual broadleaf weeds such as flixweed and stinkweed, 2,4-D should be applied to emerged, actively growing weeds in the fall the year prior to the **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** + Roundup spring pre-seeding tank-mix. Refer to the 2,4-D product label for appropriate rates.

*** Not including glyphosate tolerant canola, i.e. Roundup Ready Canola.

WEED CONTROL IN REDUCED TILLAGE FALLOW

DO NOT apply using aerial application equipment.

Treatment Notes

- 1 Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mixes in the spring to fallow land when seedling weeds have emerged, and are actively growing at the 2 to 4-leaf stage.
- 2 Reduced control may occur if applications are made at an advanced stage of weed development.

Application Directions

Weeds Controlled	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Rate	Tank Mix
Canada thistle (top growth) cow cockle flixweed green smartweed kochia lady's-thumb lamb's-quarters perennial sow-thistle (top growth) redroot pigweed Russian thistle shepherd's-purse stinkweed Tartary buckwheat wild buckwheat wild mustard	315 – 398 mL/ha +	1.1 L/ha of 2,4-D amine 500 OR 920 mL/ha of 2,4-D L.V. ester 600 in 50-100 L of water
cow cockle flixweed* <i>foxtail</i> barley** green foxtail kochia lady's-thumb lamb's-quarters redroot pigweed** Russian thistle stinkweed volunteer canola*** volunteer cereals <i>wild</i> buckwheat** wild mustard wild oats	398 mL/ha +	270-362 g ae/ha glyphosate+ 350 mL of a non- ionic surfactant registered for this use in 50-100 L of water
wild buckwheat	823 mL/ha +	270-362 g ae/ha glyphosate+ 350 mL of an approved non-ionic surfactant in 50-100 L of water

RESTRICTED

- * For control of flixweed use 362 g ae/ha of glyphosate.
- ** Suppression only.
- *** Not including glyphosate tolerant canola, i.e. Roundup Ready Canola.

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE / Roundup brand agricultural products (glyphosate only) Application Notes

- 1 These tank-mixes should be applied to emerged, actively growing annual weeds from 8-15 cm in height.
- 2 Use the higher rate of Roundup brand agricultural products (glyphosate only) when weeds are at a more advanced stage of growth.
- 3 For perennial weed control, refer to the appropriate section of this label for proper stages of growth and recommended stages of application.
- 4 Reduced control may occur if muddy water is used, such as water from dug-outs, ponds and unlined ditches.

PERENNIAL WEED CONTROL IN SUMMERFALLOW AND STUBBLE

DO NOT apply using aerial application equipment.

Treatment Notes

- 1 Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** in minimum 100 litres of water/ha.
- 2 For the most effective control of Canada thistle, follow a long-term approach that includes in crop, post-harvest, and summerfallow treatments, in conjunction with tillage operations.
- 3 If application is made after September 1st, or if soil moisture levels are extremely low after application, crop injury may occur in the spring following application.

Weeds Controlled

Weeds Controlled	Rate	Recropping in Year Following
Canada thistle curled dock (top growth) English daisy field bindweed goldenrod perennial sow thistle tansy ragwort	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone at 3.43 L/ha	cereals field corn soybeans sweet corn white beans
Canada thistle perennial sow-thistle	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE at 1.71 L/ha + 610 g ae/ha glyphosate + 350 mL of a non-ionic surfactant	<i>All of the above plus: canola</i>

Application Directions

Summerfallow Treatment Notes

- 1 Cultivate in the spring and apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** when:

Weed	Weed Stage
thistles	the majority of thistles are up and before the early bud stage (15-25 cm tall)
field bindweed	in the flowering stage
other weeds	in the early bud stage of growth

2. Cultivate three weeks after application

Stubble Treatment Notes

Apply to regrowth after harvest and at least 2 weeks prior to a killing frost.

DO NOT permit lactating dairy animals to graze fields within 7 days after application.

DO NOT harvest forage or cut hay within 30 days after application.

Withdraw meat animals from treated fields at least 3 days before slaughter.

PERENNIAL ROSETTE CONTROL IN SUMMERFALLOW

DO NOT apply using aerial application equipment.

Treatment Notes

1. For the most effective control of Canada thistle, follow a long-term approach that includes in crop, post-harvest, and summerfallow treatments, in conjunction with tillage operations.
2. Commence early spring cultivation and continue as required throughout the summer. Note: The final cultivation must occur by the end of July between July 15-August 1 and the final cultivation should cut the thistle off 5 to 7.5 cm below the soil surface.
3. Spray in minimum 100 L of water/ha when the majority of thistles have emerged as low growing rosettes 15 to 25 cm across.
4. Apply at least two weeks prior to a killing frost.
5. Cultivate three weeks after application.

Weeds Controlled

Weeds Controlled	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Rate	Recropping in Year Following
Canada thistle	1.71 L/ha	canola cereals field corn soybeans white beans

PASTURES, RANGELAND, AND NON-CROP AREAS

XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide may be used to control deciduous brush species and broadleaf weeds in non-cropland areas, such as roadsides, hydro, pipeline and railway rights-of-way, airports, military bases, wasteland and similar non-crop land areas, as well as pasture and rangeland.

Treatment Notes

For high volume handwand applications, applicators must limit volume of solution used per day to 400 L (broadleaf control spot treatment only).

For Broadleaf Weed Control

- 1 Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** or **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** tank-mixes in 110-220 L of water/ha when weeds are actively growing. Thorough coverage of foliage is necessary to control weeds.
- 2 Do not apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** or **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** tank-mixes if pasture is underseeded to legumes.

DO NOT apply using aerial application equipment.

Weeds Controlled	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Rate	Tank Mix
Canada thistle curled dock (top growth) English daisy field bindweed goldenrod perennial sow-thistle tansy ragwort	Alone at 2.9 L/ha	none
diffuse knapweed goat's beard ground cherry pasture sage poverty weed	Alone at 6.3 L/ha	none

sheep sorrel thyme-leaved spurge		
poison ivy	At 2.26 L/ha +	2.2 L/ha of 2,4-D amine (500 g/L formulation) in 560 L of water/ha
Weeds listed for XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone at 2.9 L/ha plus: wild carrot plus additional weeds found on the 2,4-D amine label	At 2.9 L/ha +	2.2 L/ha of 2,4-D amine (500 g/L formulation)
Weeds listed for XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone at 2.9 L/ha plus: wild carrot plus additional weeds found on the 2,4-D ester label	At 2.9 L/ha +	1.83 L of 2,4-D L.V. ester (600 g/L formulation)

For Brush Weed Control

1. **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** is effective in controlling many deciduous brush species that are found growing along fence rows and in other areas around the farm where they may be undesirable.
2. Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY Herbicide** tank-mixes in spring or early summer to deciduous species (leaves should be fully expanded) either as a leaf stem treatment or as a broadcast ground application.
3. Brush and trees over 2 meters tall should be cut and regrowth treated when it develops.
4. Do not apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mixes if pasture or rangeland is underseeded to legumes.
5. For Stem Foliage Treatment, apply to all foliage and stems to the point of runoff. The volume of spray mix applied per hectare will vary according to the height and density of the woody species present.
6. For Broadcast Ground Treatment, apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mixes in sufficient dilution to wet all foliage. Normally, 220-230 litres of water/ha is recommended for brush stands.

DO NOT apply using aerial application equipment.

Weeds Controlled	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Rate	Tank Mix
Alder aspen poplar cherry western snowberry (buckbrush) wolf willow (silverwillow) wild rose	At 2.9 L /1000 L of water +	4.0 L of 2,4-D amine (500 g/L formulation) OR 3.3 L of 2,4-D L.V. (600 g/L formulation)
aspen poplar	At 4.5 L/ha +	4.4 L/ha of 2,4-D amine (500 g/L formulation) OR 3.75 L/ha of 2,4-D L.V. ester (600 g/L formulation)
prickly rose	At 5.01 L/ha +	4.4 L/ha of 2,4-D amine (500 g/L formulation) OR 3.75 L/ha of 2,4-D L.V. ester (600 g/L formulation)
western snowberry	At 5.01 L/ha +	3.75 L/ha of 2,4-D L.V. ester (600 g/L formulation)

Grazing Restrictions

DO NOT permit lactating dairy animals to graze fields within 7 days after application.

DO NOT harvest forage or cut hay within 30 days after application.

Withdraw meat animals from treated fields at least 3 days before slaughter.

SEED PRODUCTION

DO NOT apply using aerial application equipment.

Treatment Notes For New/Established Stands of Red Fescue

1. Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** or **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mixes in at least 110 litres of water/ha.
2. Applications to new seedling stands may be made when the crop is 5 cm tall.
3. Application to established stands may be made up to the shot-blade stage of the crop.

4. For dandelion control, apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** plus 2,4-D amine in the fall when weeds are in the rosette or early bud stage.

Weeds Controlled	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Rate	Tank Mix
Canada thistle (top growth) clover corn spurry cow cockle green smartweed lady's-thumb perennial sow-thistle (top growth) Tartary buckwheat wild buckwheat	Alone at 823 mL/ha	none
<i>All of the above plus:</i> additional weeds found on the 2,4-D amine label	At 823 mL/ha +	1.5 L/ha of 2,4-D amine (500 g/L formulation)

For Canary seed (*Phalaris canariensis*)

1. The canary seed (*Phalaris canariensis*) should only be used as bird seed.
2. For specific weeds controlled, refer to the **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** + MCPA amine weed spectrum list under "Cereals".

Herbicide	Rate	Canary Seed (<i>Phalaris canariensis</i>) Stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone	398 mL/ha	3 - 5 leaf stage
XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE + MCPA amine	398 mL/ha + 850 mL/ha (500 g/L formulation)	3 - 5 leaf stage

For Seedling Grasses (seeded alone or underseeded with cereals)

For seed and forage production of the following seedling grasses

bromegrass, smooth	wheatgrass, crested
fescue, meadow	wheatgrass, Intermediate
fescue, tall	wheatgrass, pubescent
foxtail, meadow	wheatgrass, slender
orchard grass	wheatgrass, streambank
red fescue, creeping	wheatgrass, tall

1. Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** or **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** + tank-mixes in at least 110 litres of water/ha.
2. Application to new seedling grasses may be made when they are in the 2 to 4-leaf stage. If the seedling grass is under seeded with a cereal crop, refer to “Cereals” for additional restrictions pertaining to application type and rate.
3. If the crops are to be used as feed or pasture following treatment with XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE, **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** plus 2,4-D amine or MCPA, refer to “Grazing Restrictions”.

Weeds Controlled	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Rate	Tank Mix
Canada thistle (top growth) cleavers (higher rate only) corn spurry cow cockle green smartweed lady’s-thumb perennial sow-thistle (top growth) Tartary buckwheat wild buckwheat	Alone at 315 -398 mL/ha	none
All of the above plus: burdock (young seedlings) volunteer canola* cocklebur flixweed hemp-nettle** kochia redroot pigweed Russian pigweed Russian thistle shepherd’s-purse	At 315 - 398 mL/ha +	850 mL/ha of 2,4-D amine (500 g/L formulation) OR 850 mL/ha of MCPA amine (500 g/L formulation) OR 1.1 L/ha of MCPA K (400 g/L formulation)

volunteer sunflower*** wild radish		
---------------------------------------	--	--

* Best results will be obtained if application is made prior to bolting of canola, when this weed is at the 2 to 4 leaf stage.

** Use **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** + MCPA K for hemp-nettle control. Apply at the 2 to 3 leaf stage of weed for best control. Hemp-nettle may not be controlled if application is made at a more advanced stage of crops and weeds.

*** Depending on the growing conditions, control may be delayed slightly.

For Established Grass Pasture

- 1 Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** at 823 mL/ha with 1.5 L/ha of 2,4-D amine (500 g/L formulation) to suppress volunteer alfalfa.
- 2 Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** + 2,4-D amine in 110-220 L/ha in the spring to actively growing alfalfa at greater than 5 cm in height.

LOW-BUSH BLUEBERRIES

DO NOT apply using aerial application equipment.

Treatment Notes

1. **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** can be used alone or in a tank-mix with 2,4-D L.V. ester.
2. Apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** or the **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mix in 550 litres of water per hectare.
3. Apply in the fall while the sweet-fern is still moderately green after 90% of the blueberries have dropped their leaves. This should be done before the area is burned. Fall burning or cutting should be carried out 4 to 5 weeks after spraying. If spring burning or cutting is planned, it should be done as early as possible in the spring to reduce injury to the blueberries.

Weeds Controlled

Weeds Controlled	XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE Rate	Tank Mix
lambkill (sheep laurel) sweet fern	6.3 - 9.7 L/ha	none
additional broadleaf control	3.2 L/ha +	5.7 L of 2,4-D L.V. ester (600 g/L formulation)

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** or other Group 4 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Bayer CropScience at 1-888-283-6847 or at www.cropscience.bayer.ca.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Thaw if frozen. Shake before use.

During mixing, loading, application, clean-up and repair, workers must wear long-sleeved shirt and long pants, chemical-resistant gloves and shoes plus socks. Chemical-resistant gloves are not required inside a closed cab or closed cockpit during application. For applications to non-crop areas, applicators must also wear coveralls.

RESTRICTED

DO NOT enter treated fields until 12 hours after application to Roundup Ready 2 Xtend Soybeans and XtendFlex™ Soybeans, barley, low bush blueberries, canary seed (*Phalaris canariensis*), corn (field and sweet), fallow, oats, pastures, red fescue, spring rye, seedling grasses, stubble fields, summer fallow and wheat (spring, durum) and Roundup Ready 2 Xtend soybeans and XtendFlex™ Soybeans.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

FIRST AID

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For first aid instructions or the toxicological information essential for treatment, obtain and read the approved label from the registrant or phone the number indicated on this container.

TOXICOLOGICAL INFORMATION

Dicamba may cause severe irritation to the eyes and irritation to the skin and mucous membranes. Symptoms of overexposure to dicamba may include dizziness, muscle weakness, loss of appetite, weight loss, vomiting, decreased heart rate, shortness of breath, excitement, tenseness, depression, incontinence, cyanosis, muscle spasms, exhaustion and loss of voice.

Treat symptomatically.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and it is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

- 1 Triple- or pressure-rinse the empty container. Add the rinsings to spray mixture in the tank.
- 2 Make the empty, rinsed container unsuitable for further use.

RESTRICTED

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CLEANING SPRAY EQUIPMENT

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE alone or with 2,4-D or MCPA

If you have used **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** alone or **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** in a tank-mix with 2,4-D or MCPA, to clean the spray equipment follow these steps:

1. Thoroughly hose down the inside and outside of equipment surfaces while filling the spray tank half-full with water. Flush by operating the sprayer until the system is purged of the rinse water.
2. Fill the tank with water, adding 1 L of household ammonia for every 100 L of water. Operate the spray pump to circulate the ammonia solution through the sprayer solution for 15-20 minutes and discharge a small amount of the ammonia solution through the spray boom and nozzles.
3. Flush the solution out of the spray tank through the boom.
4. Remove the nozzles and screens and flush the system with two tanks full of water.

XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE with other Herbicides

To clean spray equipment used to apply **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** as a tank-mix with wettable powders (WP), emulsifiable concentrates (EC) or other types of water-dispersible formulations, follow these steps: (Note that if you use **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** tank-mixes with water-dispersible formulation, you must add detergent to the rinse water.)

1. Thoroughly hose down the inside and outside of equipment surfaces while filling the spray tank half-full with water. Flush by operating the sprayer until the system is purged of the rinse water.
2. Fill tank with water while adding 1 kg of detergent for every 150 litres of water. Operate the pump to circulate the detergent solution through the sprayer system for 5-10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
3. Flush the detergent solution out of the spray tank through the boom.
4. Repeat step 1 and follow steps 2 and 3.

Bulk Container Refilling

1. The container is to be refilled only with **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE**.
2. Reseal and return to an authorized Monsanto bulk site.
3. Prior to refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices.
4. Check for leaks after refilling and before transportation.
5. Do not refill or transport damaged or leaking containers.

6. For disposal, this container may be returned to the point of purchase (dealer/distributor). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.
7. If the container is not being refilled, refer to Section on "Disposal".

STORAGE

1. Store **XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE** in its original container only, away from other pesticides, fertilizer, food, or feed.
2. Keep the container closed to prevent spills and contamination.
3. Keep packages dry at all times.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Roundup Ready 2 Xtend™, Roundup Ready®, Roundup Xtend™, Roundup® and VaporGrip™ are trademarks of ©2020 Bayer Group. All rights reserved